



# Article Qubits, unitary operators, and rotations

# François Dubeau<sup>1,\*</sup>

- <sup>1</sup> Département de mathématiques Université de Sherbrooke Sherbrooke (Qc), Canada, J1K 2R1
- \* Corresponding author. François Dubeau E-mail: francois.Dubeau@usherbrooke.ca
- Academic Editor: Waqas Nazeer

Received: 10 March 2025; Accepted: 21 March 2025; Published: 23 April 2025

Abstract: We present an introduction to the mathematics of quantum physics and quantum computation which put emphasis on the basic mathematical aspects of definition and operations on qubits. We start by a comprehensive introduction of a qubit as a unit element of  $\mathbb{C}^2$ , and its representations on spheres in  $\mathbb{R}^3$ . This introduction leads to the interpretation of Pauli operators as basic rotations in  $\mathbb{R}^3$ . Then we study unitary operators. Their link to rotations in  $\mathbb{R}^3$  is established using the density operator associated to a qubit. We complete this paper by some decomposition, or splitting, problems of unitary operators on  $\mathbb{C}^2$  based on decomposition results of rotations in  $\mathbb{R}^3$ . These decomposition results are useful for the construction of quantum gates.

Keywords: Qubit, Bloch sphere, Pauli operator, unitary operator, rotation

MSC: Primary: 81P16, 81-01; Secondary: 46N50, 47N50.

# 1. Introduction

his paper is an introduction to the mathematics of quantum physics and quantum computation. The emphasis is put on the basic mathematical aspects of definitions and operations on qubits. Following [1–3], we start by a comprehensive introduction of a qubit. A qubit is the fundamental quantum state representing the smallest unit of quantum information containing one bit of classical information accessible by measurement. It is represented by or identified to a state vector which is a unit element of  $\mathbb{C}^2$ . We describe its representations on unit spheres in  $\mathbb{R}^3$ . These representations lead to the interpretation of Pauli operators as basic rotations in  $\mathbb{R}^3$ . Then we study unitary operators, and their link to rotations in  $\mathbb{R}^3$  is established using the density operator associated to a qubit. We obtain quite easily the expressions for the composition and the non-commutativity of two rotations. We complete this paper by some decomposition, or splitting, problems of unitary operators on  $\mathbb{C}^2$  obtained from results on rotations in  $\mathbb{R}^3$  [4,5]. This last subject is related to gates in quantum computation [6].

# 2. Complex vector space and qubit

# 2.1. Two dimensional complex vector space

Throughout this paper we use the Dirac bracket notation for elements of the two-dimensional complex vector space  $\mathbb{C}^2$  generated by two vectors

$$\mathbb{C}^2 = \operatorname{Lin} \{ |0\rangle, |1\rangle \} = \{ |q\rangle = a|0\rangle + b|1\rangle : a, b \in \mathbb{C} \}.$$

For  $|q_l\rangle = a_l |0\rangle + b_l |1\rangle$  (l = 1, 2) we use the Hermitian inner product

$$\langle q_1|q_2\rangle = a_1^*a_2\langle 0|0\rangle + a_1^*b_2\langle 0|1\rangle + b_1^*a_2\langle 1|0\rangle + b_1^*b_2\langle 1|1\rangle.$$

We suppose that

$$\langle i|j
angle = \delta_{ij} = \left\{egin{array}{ccc} 1 & ext{if} & i=j \ 0 & ext{if} & i
eq j \end{array} 
ight. ext{for} \quad i,j\in\{0,1\} \,,$$

so that the set  $\{|0\rangle, |1\rangle\}$  is an orthonormal basis for  $\mathbb{C}^2$ . We get

$$\langle q_1|q_2\rangle = a_1^*a_2 + b_1^*b_2,$$

and the length of  $|q\rangle$  is  $||q\rangle|| = \sqrt{\langle q|q\rangle} = \sqrt{|a|^2 + |b|^2}$ . A good introduction to the mathematics of finite-dimensional complex vector spaces is [7].

## 2.2. Definition of a qubit

A qubit is any unit length element of  $\mathbb{C}^2$ . It can be written as

$$|q\rangle = e^{i\psi} \left[ \cos\left(\theta/2\right) |0\rangle + \sin\left(\theta/2\right) e^{i\varphi} |1\rangle \right]$$

for  $\theta \in [0, \pi]$ ,  $\varphi \in [0, 2\pi)$ , and  $\psi \in \mathbb{R}$ . We will identify qubit up to its phase  $e^{i\psi}$ , i.e.

$$|q
angle\equiv|q'
angle ~~{
m iff}~~ \exists_{\psi\in\mathbb{R}}~|q'
angle=e^{i\psi}|q
angle,$$

so qubits are equivalence classes of unit length elements of  $\mathbb{C}^2$ . We will use

$$|q\rangle = \cos(\theta/2) |0\rangle + \sin(\theta/2) e^{i\varphi} |1\rangle,$$

as its representative.

A qubit  $|q^{\perp}\rangle$ , orthogonal to the qubit  $|q\rangle$ , is given by

$$|q^{\perp}\rangle = \cos\left(\frac{\pi-\theta}{2}\right)|0
angle + \sin\left(\frac{\pi-\theta}{2}\right)e^{i(\pi+\varphi)}|1
angle,$$

so that  $\langle q | q^{\perp} \rangle = 0$ .

# 3. Spherical representations of a qubit

### 3.1. First representation

We can represent qubits on the upper-half unit sphere in  $\mathbb{R}^3$ . Each qubit  $|q\rangle$  with  $\theta \in [0, \pi)$  is represented by a unique point, while for  $\theta = \pi$ ,  $|q\rangle = e^{i\varphi}|1\rangle \equiv |1\rangle$  is represented by the whole unit circle  $x^2 + y^2 = 1$  and z = 0.



Figure 1. Upper-half unit sphere representation

For example,  $|q\rangle$  and  $|q^{\perp}\rangle$  are indicated on Figure 1. Their corresponding vectors are

$$\begin{cases} x = \sin\left(\frac{\theta}{2}\right)\cos(\varphi), \\ y = \sin\left(\frac{\theta}{2}\right)\sin(\varphi), \\ z = \cos\left(\frac{\theta}{2}\right), \end{cases} \text{ and } \begin{cases} x^{\perp} = \sin\left(\frac{\pi-\theta}{2}\right)\cos(\pi+\varphi), \\ y^{\perp} = \sin\left(\frac{\pi-\theta}{2}\right)\sin(\pi+\varphi), \\ z^{\perp} = \cos\left(\frac{\pi-\theta}{2}\right). \end{cases}$$

## 3.2. Second representation: the Bloch sphere

To obtain only one point on the sphere for  $|1\rangle$ , we continuously transform the point (x, y, z) as follows

$$\begin{cases} x_{\lambda} &= \sin\left(\frac{1+\lambda}{2}\theta\right)\cos(\varphi), \\ y_{\lambda} &= \sin\left(\frac{1+\lambda}{2}\theta\right)\sin(\varphi), \quad \text{ for } \lambda \in [0,1], \\ z_{\lambda} &= \cos\left(\frac{1+\lambda}{2}\theta\right), \end{cases}$$

At the end of the transformation, for  $\lambda = 1$ , we get

$$\begin{cases} x_1 &= \sin(\theta)\cos(\varphi), \\ y_1 &= \sin(\theta)\sin(\varphi), \\ z_1 &= \cos(\theta). \end{cases}$$

Hence the qubit  $|q\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + \sin\left(\frac{\theta}{2}\right)e^{i\varphi}|1\rangle$  is represented by the vector

 $\vec{q} = (\sin(\theta)\cos(\varphi), \sin(\theta)\sin(\varphi), \cos(\theta)),$ 

on the unit sphere of  $\mathbb{R}^3$ . On the same way, for  $|q^{\perp}\rangle$  we get

$$\vec{q}^{\perp} = (\sin(\pi - \theta)\cos(\pi + \varphi), \sin(\pi - \theta)\sin(\pi + \varphi), \cos(\pi - \theta))$$
  
=  $-(\sin(\theta)\cos(\varphi), \sin(\theta)\sin(\varphi), \cos(\theta))$   
=  $-\vec{q}.$ 

This representation on the unit sphere of  $\mathbb{R}^3$  is called the Bloch sphere representation of a qubit (see Figure 2), and we have the next result.



Figure 2. Bloch sphere representation

**Theorem 1.** The correspondence  $|q\rangle \leftrightarrow \vec{q}$  is a bijection.

# 4. Elementary rotations in $\mathbb{R}^3$ and Pauli operators

The spherical representation for qubits in  $\mathbb{R}^3$  suggests to take a look at rotations in  $\mathbb{R}^3$ . Let us start by considering rotations around the axis *OX*, *OY*, and *OZ*. More general rotations will be considered later in this text.

### **4.1. Rotation around** *OX*

The rotation of an angle  $\xi$  around the vector  $\vec{i}$  is given by

$$R_{(\vec{i},\xi)} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\xi) & -\sin(\xi) \\ 0 & \sin(\xi) & \cos(\xi) \end{bmatrix},$$

and for  $\xi = \pi$  we get

$$\vec{q}' = R_{(\vec{i},\pi)}(\vec{q}) = \left(\begin{array}{c} \sin\left(\pi - \theta\right)\cos\left(-\varphi\right) \\ \sin\left(\pi - \theta\right)\sin\left(-\varphi\right) \\ \cos\left(\pi - \theta\right) \end{array}\right).$$

So the corresponding qubit is

$$\begin{aligned} |q'\rangle &= \cos\left(\frac{\pi-\theta}{2}\right)|0\rangle + \sin\left(\frac{\pi-\theta}{2}\right)e^{-i\varphi}|1\rangle \\ &= e^{-i\varphi}\left[\cos\left(\theta/2\right)|1\rangle + \sin\left(\theta/2\right)e^{i\varphi}|0\rangle\right]. \end{aligned}$$

If we set  $\hat{\sigma}_x |0\rangle = |1\rangle$  and  $\hat{\sigma}_x |1\rangle = |0\rangle$ , we get  $|q'\rangle = e^{-i\varphi} \hat{\sigma}_x |q\rangle \equiv \hat{\sigma}_x |q\rangle$ .

## **4.2. Rotation around** *OY*

For the rotation around the vector  $\vec{j}$  we have

$$R_{(\vec{j},\xi)} = \left[ egin{array}{ccc} \cos(\xi) & 0 & \sin(\xi) \ 0 & 1 & 0 \ -\sin(\xi) & 0 & \cos(\xi) \end{array} 
ight].$$

For  $\xi = \pi$  we get

$$\vec{q}' = R_{(\vec{j},\pi)}(\vec{q}) = \begin{pmatrix} \sin(\pi - \theta)\cos(\pi - \varphi) \\ \sin(\pi - \theta)\sin(\pi - \varphi) \\ \cos(\pi - \theta) \end{pmatrix},$$

hence

$$\begin{aligned} |q'\rangle &= \cos\left(\frac{\pi-\theta}{2}\right)|0\rangle + \sin\left(\frac{\pi-\theta}{2}\right)e^{i(\pi-\varphi)}|1\rangle \\ &= e^{i(\frac{\pi}{2}-\varphi)}\left[\cos\left(\theta/2\right)i|1\rangle + \sin\left(\theta/2\right)e^{i\varphi}(-i)|0\rangle\right]. \end{aligned}$$

If we set  $\hat{\sigma}_y |0\rangle = i|1\rangle$  and  $\hat{\sigma}_y |1\rangle = -i|0\rangle$ , we get  $|q'\rangle = e^{i(\frac{\pi}{2} - \varphi)} \hat{\sigma}_y |q\rangle \equiv \hat{\sigma}_y |q\rangle$ .

## **4.3. Rotation around** *OZ*

For the rotation around the vector  $\vec{k}$  we have

$$R_{(\vec{k},\xi)} = \begin{bmatrix} \cos(\xi) & -\sin(\xi) & 0\\ \sin(\xi) & \cos(\xi) & 0\\ 0 & 0 & 1 \end{bmatrix},$$

so

$$\vec{q}' = R_{(\vec{k},\pi)}(\vec{q}) = \left( egin{array}{c} \sin\left(\theta\right)\cos\left(\pi+\varphi
ight) \\ \sin\left(\theta\right)\sin\left(\pi+\varphi
ight) \\ \cos\left(\theta
ight) \end{array} 
ight),$$

and

$$|q'\rangle = \cos(\theta/2) |0\rangle + \sin(\theta/2) e^{i(\pi+\varphi)} |1\rangle$$
  
=  $\cos(\theta/2) |0\rangle + \sin(\theta/2) e^{i\varphi} (-1) |1\rangle.$ 

If we set  $\hat{\sigma}_z |0\rangle = |0\rangle$  and  $\hat{\sigma}_z |1\rangle = -|1\rangle$ , we get  $|q'\rangle = \hat{\sigma}_z |q\rangle$ .

#### 4.4. Pauli operators

The linear operators  $\hat{\sigma}_x$ ,  $\hat{\sigma}_y$ , and  $\hat{\sigma}_z$  defined above form the set of Pauli operators on  $\mathbb{C}^2$ . They correspond to half-revolutions in  $\mathbb{R}^3$ .

## 5. Linear operators on $\mathbb{C}^2$

Let  $\hat{A} : \mathbb{C}^2 \to \mathbb{C}^2$  be a linear operator. Its adjoint  $\hat{A}^+$  is defined by the relation

$$\langle v|\hat{A}^+|u\rangle = \langle v|(\hat{A}^+|u\rangle)\rangle = \langle u|(\hat{A}|v\rangle)\rangle^* = \langle u|\hat{A}|v\rangle^*,$$

for all  $|u\rangle$ ,  $|v\rangle \in \mathbb{C}^2$ . It is easy to observe that  $(\hat{B}\hat{A})^+ = \hat{A}^+\hat{B}^+$ . We say that  $\hat{A}$  is a positive linear operator iff (if and only if)  $\langle u|\hat{A}|u\rangle \geq 0$  for all  $|u\rangle \in \mathbb{C}^2$ .

Let us consider the following definitions:

- $\hat{A}$  is a normal operator iff  $\hat{A}\hat{A}^+ = \hat{A}^+\hat{A}$ ;
- $\hat{A}$  is a unitary operator iff  $\hat{A}^+ = \hat{A}^{-1}$ ;
- $\hat{A}$  is a Hermitian operator iff  $\hat{A}^+ = \hat{A}$ .

Obviously, any unitary or Hermitian operator is also a normal operator.

For  $|q\rangle \in \mathbb{C}^2$ , we define the external product  $|q\rangle\langle q|$  (also known as a projection operator) which is a positive Hermitian linear operator. Indeed

- positive: because  $\langle p|q\rangle\langle q|p\rangle = |\langle q|p\rangle|^2 \ge 0$ ;
- Hermitian (and hence normal) because  $(|q\rangle\langle q|)^+ = |q\rangle\langle q|$ .

We also observe that

$$\left(|q\rangle\langle q|\right)^{k} = \|q\|^{2(k-1)} |q\rangle\langle q|,$$

for k = 1, 2, 3, ...

For any orthonormal basis  $\{|v_0\rangle, |v_1\rangle\}$  of  $\mathbb{C}^2$ , we get the identity operator, noted  $\hat{\sigma}_o$ , from

$$\hat{\sigma}_o = |v_0\rangle \langle v_0| + |v_1\rangle \langle v_1|.$$

Also, using any orthonormal basis  $\{|v_0\rangle, |v_1\rangle\}$  of  $\mathbb{C}^2$ , the trace of a linear operator is given by

$$\operatorname{Tr}(\hat{A}) = \langle v_0 | \hat{A} | v_0 \rangle + \langle v_1 | \hat{A} | v_1 \rangle,$$

and this value does not depend on the choice of the orthonormal basis used for the computation.

### 6. More about Pauli operators and related bases

For two vectors  $\vec{u}$  and  $\vec{v}$  of  $\mathbb{R}^3$ , the dot and cross products are respectively noted by  $\vec{u} \cdot \vec{v}$  and  $\vec{u} \times \vec{v}$ . We recall here standard results about dot and cross products which will used in the next sections. Proofs of these results can be found for example in [8,9].

**Lemma 1.** Let  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$ , and  $\vec{d}$  in  $\mathbb{R}^3$ . We have

- $(\vec{a} \times \vec{b}) = -(\vec{b} \times \vec{a});$
- $\vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{c} \cdot (\vec{a} \times \vec{b}) = \vec{b} \cdot (\vec{c} \times \vec{a});$
- $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} (\vec{a} \cdot \vec{b})\vec{c};$
- $(\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) = (\vec{a} \cdot \vec{c})(\vec{b} \cdot \vec{d}) (\vec{b} \cdot \vec{c})(\vec{a} \cdot \vec{d});$
- $\vec{d} \cdot (\vec{a} \times (\vec{b} \times \vec{c})) = (\vec{d} \times \vec{a}) \cdot (\vec{b} \times \vec{c}).$

The set  $S = \{\hat{\sigma}_o, \hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z\}$  form a basis of the 4-dimensional vector space of linear operators on  $\mathbb{C}^2$ . Table 1 presents the composition table for the 4 elements of this basis.

	$\hat{\sigma}_o$	$\hat{\sigma}_x$	$\hat{\sigma}_y$	$\hat{\sigma}_z$
$\hat{\sigma}_o$	$\hat{\sigma}_{o}$	$\hat{\sigma}_{\chi}$	$\hat{\sigma}_y$	$\hat{\sigma}_z$
$\hat{\sigma}_{x}$	$\hat{\sigma}_x$	$\hat{\sigma}_o$	$i\hat{\sigma}_z$	$-i\hat{\sigma}_y$
$\hat{\sigma}_y$	$\hat{\sigma}_y$	$-i\hat{\sigma}_z$	$\hat{\sigma}_o$	$i\hat{\sigma}_x$
$\hat{\sigma}_z$	$\hat{\sigma}_z$	iô <sub>y</sub>	$-i\hat{\sigma}_x$	$\hat{\sigma}_o$

Table 1. Multiplication table for Pauli operators

**Proposition 1.** The Pauli operators are unitary and Hermitien, and hence normal.

Let  $\vec{v} = (v_x, v_y, v_z) \in \mathbb{R}^3$  and  $\hat{\sigma} = (\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z)$ . We define the operator

 $\vec{v} \cdot \hat{\sigma} = v_x \hat{\sigma}_x + v_y \hat{\sigma}_y + v_z \hat{\sigma}_z.$ 

Next are some basic results useful for the next sections.

**Proposition 2.** For  $\vec{a}$  and  $\vec{b}$  in  $\mathbb{R}^3$ 

$$(\vec{a}\cdot\hat{\sigma})(\vec{b}\cdot\hat{\sigma}) = (\vec{a}\cdot\vec{b})\hat{\sigma}_o + i(\vec{a}\times\vec{b})\cdot\hat{\sigma}.$$

**Corollary 1.** For  $\vec{n}$  in  $\mathbb{R}^3$ ,  $(\vec{n} \cdot \hat{\sigma})(\vec{n} \cdot \hat{\sigma}) = \|\vec{n}\|^2 \hat{\sigma}_o$ .

**Corollary 2.** For  $\vec{a}$  and  $\vec{b}$  in  $\mathbb{R}^3$ 

- $(\vec{a} \cdot \hat{\sigma})(\vec{b} \cdot \hat{\sigma}) + (\vec{b} \cdot \hat{\sigma})(\vec{a} \cdot \hat{\sigma}) = 2(\vec{a} \cdot \vec{b})\hat{\sigma}_o,$   $(\vec{a} \cdot \hat{\sigma})(\vec{b} \cdot \hat{\sigma}) (\vec{b} \cdot \hat{\sigma})(\vec{a} \cdot \hat{\sigma}) = 2i(\vec{a} \times \vec{b}) \cdot \hat{\sigma}.$

**Corollary 3.** For  $\vec{a}$  and  $\vec{b}$  in  $\mathbb{R}^3$ 

- $(\vec{a}\cdot\hat{\sigma})(\vec{b}\cdot\hat{\sigma})(\vec{a}\cdot\hat{\sigma}) = \|\vec{a}\|^2 (\vec{b}\cdot\hat{\sigma}) + 2(\vec{a}\cdot\vec{b})(\vec{a}\cdot\hat{\sigma}),$   $(\vec{a}\cdot\hat{\sigma})(\vec{b}\cdot\hat{\sigma})(\vec{a}\cdot\hat{\sigma}) = \|\vec{a}\|^2 (\vec{b}\cdot\hat{\sigma}) + 2i(\vec{a}\times\vec{b})\cdot\hat{\sigma}(\vec{a}\cdot\hat{\sigma}).$

**Proposition 3.** *For the qubit*  $|q\rangle \in \mathbb{C}^2$ 

•  $\vec{q} \cdot \hat{\sigma} |q\rangle = |q\rangle$ , •  $\vec{q} \cdot \hat{\sigma} |q^{\perp}\rangle = -|q^{\perp}\rangle$ .

**Proposition 4.** *For the qubit*  $|q\rangle \in \mathbb{C}^2$ ,  $\langle q|\hat{\sigma}|q\rangle = \vec{q}$ .

**Proposition 5.**  $\operatorname{Tr}(\hat{\sigma}_0) = 2$ , and  $\operatorname{Tr}(\hat{\sigma}_l) = 0$  for l = x, y, z.

**Proposition 6.** *For any*  $\vec{n} \in \mathbb{R}^3$ *, let* 

$$(\vec{n}\cdot\hat{\sigma})\hat{\sigma} = \left((\vec{n}\cdot\hat{\sigma})\hat{\sigma}_x, (\vec{n}\cdot\hat{\sigma})\hat{\sigma}_y, (\vec{n}\cdot\hat{\sigma})\hat{\sigma}_z\right),$$

then

$$\operatorname{Tr}((\vec{n}\cdot\hat{\sigma})\hat{\sigma})=\vec{n}.$$

**Proposition 7.** We have

$$\begin{cases} \hat{\sigma}_x \hat{\sigma}_y \hat{\sigma}_z &= \hat{\sigma}_y \hat{\sigma}_z \hat{\sigma}_x &= \hat{\sigma}_z \hat{\sigma}_x \hat{\sigma}_y &= i\hat{\sigma}_o &= e^{i\frac{\pi}{2}}, \\ \hat{\sigma}_z \hat{\sigma}_y \hat{\sigma}_x &= \hat{\sigma}_y \hat{\sigma}_x \hat{\sigma}_z &= \hat{\sigma}_x \hat{\sigma}_z \hat{\sigma}_y &= -i\hat{\sigma}_o &= e^{-i\frac{\pi}{2}}. \end{cases}$$

To get some results later in this text, we introduce a second basis for the space of linear operators on  $\mathbb{C}^2$ 

$$\mathcal{P}=\left\{ egin{array}{l} \hat{\pi}_{ij}=ert i
angle \langle jert \,:\,\,i,j=0,1 
ight. 
ight\}$$
 ,

where  $\hat{\pi}_{ii}$  is a projection for i = j and a permutation for  $i \neq j$ . The two bases are related by the formula

$$\begin{cases} \hat{\pi}_{00} &= \frac{1}{2} \left( \hat{\sigma}_{o} + \hat{\sigma}_{z} \right), \\ \hat{\pi}_{01} &= \frac{1}{2} \left( \hat{\sigma}_{x} + i \hat{\sigma}_{y} \right), \\ \hat{\pi}_{10} &= \frac{1}{2} \left( \hat{\sigma}_{x} - i \hat{\sigma}_{y} \right), \\ \hat{\pi}_{11} &= \frac{1}{2} \left( \hat{\sigma}_{o} - \hat{\sigma}_{z} \right), \end{cases} \text{ and } \begin{cases} \hat{\sigma}_{o} &= \hat{\pi}_{00} + \hat{\pi}_{11}, \\ \hat{\sigma}_{x} &= \hat{\pi}_{10} + \hat{\pi}_{01}, \\ \hat{\sigma}_{y} &= i \left[ \hat{\pi}_{10} - \hat{\pi}_{01} \right] \\ \hat{\sigma}_{z} &= \hat{\pi}_{00} - \hat{\pi}_{11}. \end{cases}$$

We get

$$\begin{cases} \hat{\pi}_{00}^+ &= \hat{\pi}_{00} \\ \hat{\pi}_{11}^+ &= \hat{\pi}_{11} \end{cases} \text{ and } \begin{cases} \hat{\pi}_{01}^+ &= \hat{\pi}_{10}, \\ \hat{\pi}_{10}^+ &= \hat{\pi}_{01}. \end{cases}$$

### 7. Unitary operator

Any linear operator  $\hat{U}$  can be expressed as a linear combination of the Pauli operators

$$\hat{U}=e^{iarphi}\left[r_{0}\hat{\sigma}_{0}+ec{r}\cdot\hat{\sigma}
ight]$$
 ,

with a real number  $r_0 \ge 0$ , and  $\vec{r} \in \mathbb{C}^3$ . From the normality condition  $\hat{U}U = U\hat{U}$  we get  $\vec{r} = \lambda \vec{n}$  where  $\vec{n}$  is a unit vector in  $\mathbb{R}^3$  and  $\lambda \in \mathbb{C}$ . From the unitary condition  $\hat{U}U = \hat{\sigma}_0 = U\hat{U}$  we get

$$\begin{cases} r_0^2 + |\lambda|^2 &= 1, \\ r_0 (\lambda + \lambda^*) &= 0. \end{cases}$$

So  $\lambda$  is a purely imaginary number,  $\lambda = i\tilde{\lambda}$  with  $\tilde{\lambda} \in \mathbb{R}$ . Let us set

$$\begin{cases} r_0 = \cos(\xi/2), & \text{for} \quad \xi \in [0,\pi].\\ |\tilde{\lambda}| = \sin(\xi/2), & \end{cases}$$

It follows that any unitary operator  $\hat{U}$  can be expressed as

$$\hat{U} = e^{i\varphi} \left[ \cos(\xi/2)\hat{\sigma}_0 - i\sin(\xi/2)\vec{n}\cdot\hat{\sigma} \right]$$

with  $\xi \in [0, \pi]$ , and  $\vec{n} \in \mathbb{R}^3$  such that  $\|\vec{n}\| = 1$ .

### 8. The density operator

The density operator associated to the qubit  $|q\rangle = \cos(\theta/2) |0\rangle + \sin(\theta/2) e^{i\varphi} |1\rangle$  is the external product

$$\hat{\rho} = |q\rangle\langle q|.$$

Using the link between the two bases S and P, we get

$$\hat{\rho} = rac{1}{2} \left[ \hat{\sigma}_0 + \vec{q} \cdot \hat{\sigma} 
ight].$$

**Theorem 2.** The density operator  $\hat{\rho}$  associated to the qubit  $|q\rangle \in \mathbb{C}^2$  is a positive Hermitian linear operator such that  $\operatorname{Tr}(\hat{\rho}) = 1$  and  $\hat{\rho}^2 = \hat{\rho}$ .

**Proof.** Direct consequence of the fact that  $\hat{\rho} = |q\rangle\langle q|$  is an external product.  $\Box$ 

**Theorem 3.** If  $\hat{\rho}$  is a positive Hermitian linear operator such that  $\operatorname{Tr}(\hat{\rho}) = 1$  and  $\hat{\rho}^2 = \hat{\rho}$ , then there exits a qubit  $|q\rangle \in \mathbb{C}^2$  such that  $\hat{\rho} = |q\rangle\langle q|$ .

**Proof.** If  $\hat{\rho}$  is normal operator, then  $\hat{\rho} = r_0 \hat{\sigma}_0 + \lambda \vec{v} \cdot \hat{\sigma}$  with  $r_0 \ge 0$  and  $\vec{v}$  is a unit vector in  $\mathbb{R}^3$ . Since  $\hat{\rho}$  is Hermitian, it follows that  $\lambda \in \mathbb{R}$ . Since  $\text{Tr}(\hat{\rho}) = 2r_0$ , then  $r_0 = 1/2$ . For any qubit  $|p\rangle$  we have

$$\langle p|\hat{
ho}|p\rangle = \frac{1}{2}\langle p|\hat{\sigma}_0|p\rangle + \lambda \vec{v} \cdot \langle p|\hat{\sigma}|p\rangle$$

$$= \frac{1}{2} + \lambda \vec{v} \cdot \vec{p},$$

because  $\langle p | \hat{\sigma}_0 | p \rangle = \langle p | p \rangle = 1$  and  $\langle p | \hat{\sigma} | p \rangle = \vec{p}$ . So, for any  $\vec{p}$  we have

$$\langle p|\hat{
ho}|p
angle \geq rac{1}{2} - |\lambda|.$$

But

$$\langle p|\hat{
ho}|p
angle = rac{1}{2} - |\lambda|,$$

for  $\vec{p} = \vec{v}$  or  $\vec{p} = -\vec{v}$ . For positivity we must have  $\frac{1}{2} - |\lambda| \ge 0$ , so  $|\lambda| \le \frac{1}{2}$ . With the last condition we have

$$\hat{
ho}^2 = \left(rac{1}{4} + \lambda^2
ight)\hat{\sigma_0} + \lambda\hat{artheta}\cdot\hat{\sigma}$$

and  $\hat{\rho}^2 = \hat{\rho}$  iff  $\frac{1}{4} + \lambda^2 = \frac{1}{2}$ , hence  $|\lambda| = \frac{1}{2}$ . Let us set

$$\vec{q} = \begin{cases} \vec{v} & \text{if } \lambda = 1/2, \\ -\vec{v} & \text{if } \lambda = -1/2, \end{cases}$$

so we get

$$\hat{
ho} = rac{1}{2} \left[ \hat{\sigma}_0 + ec{q} \cdot \hat{\sigma} 
ight]$$
 ,

for a unit vector  $\vec{q} \in \mathbb{R}^3$ . But there exist  $\theta \in [0, \pi]$  and  $\varphi \in [0, 2\pi)$  such that

 $\vec{q} = (\sin(\theta)\cos(\varphi), \sin(\theta)\sin(\varphi), \cos(\theta)),$ 

which is associated to the qubit

 $|q\rangle = \cos(\theta/2) |0\rangle + \sin(\theta/2) e^{i\varphi} |1\rangle.$ 

It follows that  $\hat{\rho}$  is the density operator associated to the qubit  $|q\rangle$ .  $\Box$ 

The correspondences  $|q\rangle \leftrightarrow \vec{q} \leftrightarrow \hat{\rho}$  are bijections, which means that  $|q\rangle$  determine uniquely  $\hat{\rho}$  and  $\hat{\rho}$ determine uniquely  $|q\rangle$ .

### 9. Rotations on qubits

## 9.1. Definition

If we identify unitary operator up to its phase, a rotation in  $\mathbb{C}^2$  is given by

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \cos\left(\xi/2\right)\hat{\sigma}_0 - i\sin\left(\xi/2\right)\vec{n}\cdot\hat{\sigma},$$

where the angle  $\xi \in [0, \pi]$  and the unit vector  $\vec{n} \in \mathbb{R}^3$  are uniquely determined.

## 9.2. Basic properties of rotations

We start by mentioning some basic properties of rotations.

Proposition 8. We can verify directly that

- $\widehat{\operatorname{rot}}_{\vec{n}}(0) = \hat{\sigma}_o$ ,

- $\operatorname{rot}_{\vec{n}}(0) = v_o$ ,  $\operatorname{rot}_{\vec{n}}(\xi_1) \operatorname{rot}_{\vec{n}}(\xi_2) = \operatorname{rot}_{\vec{n}}(\xi_1 + \xi_2)$ ,  $\operatorname{rot}_{-\vec{n}}(\xi) = \operatorname{rot}_{\vec{n}}(-\xi)$ ,  $\operatorname{rot}_{\vec{n}}^{-1}(\xi) = \operatorname{rot}_{\vec{n}}(-\xi)$ ,  $\operatorname{rot}_{\vec{n}}^+(\xi) = \operatorname{rot}_{\vec{n}}^{-1}(\xi)$ ,  $\|\operatorname{rot}_{\vec{n}}(\xi)|q\rangle\| = \||q\rangle\|$  for any  $|q\rangle \in \mathbb{C}^2$ .

# 9.3. Link between rotations on $\mathbb{C}^2$ and $\mathbb{R}^3$

For a qubit  $|q\rangle$ , let us set

$$|q'\rangle = \operatorname{rot}_{\vec{n}}(\xi)|q\rangle.$$

The vector  $\vec{q}'$  can be identified from the density operator  $\hat{\rho}'$  since

$$\hat{
ho}' = |q'
angle \langle q'| = rac{1}{2} \left[\hat{\sigma}_0 + \vec{q}\;'\cdot\hat{\sigma}
ight]$$

But

$$\hat{\rho}' = \widehat{\operatorname{rot}}_{\vec{n}}(\xi) |q\rangle \langle q| \widehat{\operatorname{rot}}_{\vec{n}}^+(\xi) = \cos^2(\xi/2)\hat{\rho} + i\sin(\xi/2)\cos(\xi/2) \left[\hat{\rho}\left(\vec{n}\cdot\hat{\sigma}\right) - \left(\vec{n}\cdot\hat{\sigma}\right)\hat{\rho}\right] + \sin^2(\xi/2)\left(\vec{n}\cdot\hat{\sigma}\right)\hat{\rho}\left(\vec{n}\cdot\hat{\sigma}\right)$$

Let us introduce the following lemma to simplify the computation.

**Lemma 2.** For any qubit  $|q\rangle \in \mathbb{C}^2$  and  $\vec{n} \in \mathbb{R}^3$ , we have

- $\hat{\rho}(\vec{n}\cdot\hat{\sigma}) = \frac{1}{2} [(\vec{q}\cdot\vec{n})\hat{\sigma}_0 + (\vec{n}+i(\vec{q}\times\vec{n}))\cdot\hat{\sigma}],$   $(\vec{n}\cdot\hat{\sigma})\hat{\rho} = \frac{1}{2} [(\vec{n}\cdot\vec{q})\hat{\sigma}_0 + (\vec{n}+i(\vec{n}\times\vec{q}))\cdot\hat{\sigma}],$   $(\vec{n}\cdot\hat{\sigma})\hat{\rho}(\vec{n}\cdot\hat{\sigma}) = \frac{1}{2} [\hat{\sigma}_0 + (2(\vec{q}\cdot\vec{n})\vec{n}-\vec{q})\cdot\hat{\sigma}].$

So, using this lemma we get

$$\hat{
ho}' = rac{1}{2} \left[ \hat{\sigma}_0 + \mathrm{R}_{(\vec{n}, \xi)}(\vec{q}) \cdot \hat{\sigma} 
ight]$$
 ,

where

$$\vec{q}' = \mathrm{R}_{(\vec{n},\xi)}(\vec{q}) = \cos(\xi)\vec{q} + \sin(\xi)\left(\vec{n}\times\vec{q}\right) + \left(1-\cos(\xi)\right)\left(\vec{n}\cdot\vec{q}\right)\vec{n}$$

is the finite rotation formula in  $\mathbb{R}^3$ . This expression is often called the Rodrigues' rotation formula, see [3,5] for example, but it would be more appropriate to name it Euler's rotation formula because it is Euler who was first to derive the finite rotation formula [10].

Consequently, applying  $\widehat{\text{rot}}_{\vec{n}}(\xi)$  on  $|q\rangle$  is equivalent to applying  $R_{(\vec{n},\xi)}$  on  $\vec{q}$ , in other words we have

$$|q'\rangle = \widehat{\operatorname{rot}}_{\vec{n}}(\xi)|q\rangle$$
 iff  $\vec{q}' = \operatorname{R}_{(\vec{n},\xi)}(\vec{q}).$ 

If we set

- $\hat{\mathbf{n}}^{0}(\vec{q}) = \vec{q}$ ,  $\hat{\mathbf{n}}^{1}(\vec{q}) = \vec{n} \times \vec{q}$ ,  $\hat{\mathbf{n}}^{2}(\vec{q}) = \vec{n} \times (\vec{n} \times \vec{q}) = (\vec{n} \cdot \vec{q})\vec{n} \vec{q}$ ,

we can rewrite the finite rotation formula as

$$\mathbf{R}_{(\vec{n},\xi)} = \hat{\mathbf{n}}^0 + \sin(\xi)\hat{\mathbf{n}}^1 + (1 - \cos(\xi))\hat{\mathbf{n}}^2.$$

#### 10. Operations on rotations

#### 10.1. Composition of rotations

Let us look at the composition of two rotations  $\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1)$  and  $\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)$ . We have

$$\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1) = X\hat{\sigma}_0 - i\vec{N}\cdot\hat{\sigma}_1$$

where

$$\begin{cases} X = \cos(\xi_2/2)\cos(\xi_1/2) - \sin(\xi_2/2)\sin(\xi_1/2)\vec{n}_2 \cdot \vec{n}_1, \\ \vec{N} = \cos(\xi_1/2)\sin(\xi_2/2)\vec{n}_2 + \cos(\xi_2/2)\sin(\xi_1/2)\vec{n}_1 + \sin(\xi_2/2)\sin(\xi_1/2)\vec{n}_2 \times \vec{n}_1. \end{cases}$$

Since

$$|X|^2 + \left\|\vec{N}\right\|^2 = 1,$$

we choose  $\xi_0 \in [0, 2\pi]$  such that  $X = \cos(\xi_0/2)$  and set  $\vec{N} = \sin(\xi_0/2)\vec{n}_0$  to get

$$\operatorname{rot}_{\vec{n}_2}(\xi_2)\operatorname{rot}_{\vec{n}_1}(\xi_1) = \cos(\xi_0/2)\hat{\sigma}_0 - i\sin(\xi_0/2)\vec{n}_0 \cdot \hat{\sigma} = \operatorname{rot}_{\vec{n}_0}(\xi_0)$$

with

Ś

$$\begin{cases} \cos(\xi_0/2) &= \cos(\xi_2/2)\cos(\xi_1/2) - \sin(\xi_2/2)\sin(\xi_1/2)\vec{n}_2 \cdot \vec{n}_1, \\ \sin(\xi_0/2)\vec{n}_0 &= \cos(\xi_1/2)\sin(\xi_2/2)\vec{n}_2 + \cos(\xi_2/2)\sin(\xi_1/2)\vec{n}_1 + \sin(\xi_2/2)\sin(\xi_1/2)\vec{n}_2 \times \vec{n}_1. \end{cases}$$

For rotations in  $\mathbb{R}^3$ , we consider the density operators to get

$$\frac{1}{2} \left[ \hat{\sigma}_0 + \mathbf{R}_{(\vec{n}_0, \xi_0)}(\vec{q}) \cdot \hat{\sigma} \right] = \frac{1}{2} \left[ \hat{\sigma}_0 + \mathbf{R}_{(\vec{n}_2, \xi_2)} \left( \mathbf{R}_{(\vec{n}_1, \xi_1)}(\vec{q}) \right) \cdot \hat{\sigma} \right].$$

So

$$\mathbf{R}_{(\vec{n}_2,\xi_2)}\mathbf{R}_{(\vec{n}_1,\xi_1)} = \mathbf{R}_{(\vec{n}_0,\xi_0)}.$$

There are many ways to get this result [11], but following [10] it was first established by Rodrigues. Since

$$\begin{array}{lll} \sin(\xi_0) \hat{n}_0^1 & = & 2\cos(\xi_0/2)\sin(\xi_0/2)\hat{n}_0^1, \\ (1 - \cos(\xi_0)) \hat{n}_0^2 & = & 2\left(\sin(\xi_0/2)\hat{n}_0^1\right)^2, \end{array}$$

we can evaluate

$$\mathbf{R}_{(\vec{n}_i,\xi_i)} = \hat{\mathbf{n}}_i^0 + \sin(\xi_i)\hat{\mathbf{n}}_i^1 + (1 - \cos(\xi_i))\hat{\mathbf{n}}_i^2$$

not only for i = 1, 2 but also for i = 0.

Let us remark that for  $\vec{n}_1 = \vec{n}_2 = \vec{n}$  we get

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}}(\xi_1) = \widehat{\operatorname{rot}}_{\vec{n}}(\xi_1 + \xi_2).$$

and

$$R_{(\vec{n},\xi_1)}R_{(\vec{n},\xi_2)} = R_{(\vec{n},\xi_1+\xi_2)}.$$

### 10.2. Non-commutativity of rotations

From the last remark, the composition of two rotations around the same axis is commutative. The non-commutativity for the composition of rotations around different axes of rotation, illustrated for example in [12], is characterized by the next result.

#### Theorem 4. We have

$$\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1) = \widehat{\operatorname{rot}}_{R_{(\vec{n}_2,\xi_2)}(\vec{n}_1)}(\xi_1)\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)$$

or

$$\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1)\widehat{\operatorname{rot}}_{\vec{n}_2}(-\xi_2) = \widehat{\operatorname{rot}}_{R_{(\vec{n}_2,\xi_2)}(\vec{n}_1)}(\xi_1).$$

**Proof.** We have to show that  $\widehat{\operatorname{rot}}_{\vec{n}_0}(\xi_0) = \widehat{\operatorname{rot}}_{\vec{n}'_0}(\xi'_0)$  where

$$\begin{array}{lll} & (\widehat{\operatorname{rot}}_{\vec{n}_{2}}(\xi_{2})\widehat{\operatorname{rot}}_{\vec{n}_{1}}(\xi_{1}) & = & \widehat{\operatorname{rot}}_{\vec{n}_{0}}(\xi_{0}), \\ & & (\widehat{\operatorname{rot}}_{R_{(\vec{n}_{2},\vec{\xi}_{2})}(\vec{n}_{1})}(\xi_{1})\widehat{\operatorname{rot}}_{\vec{n}_{2}}(\xi_{2}) & = & \widehat{\operatorname{rot}}_{\vec{n}_{0}}(\xi_{0}'). \end{array}$$

But

$$\begin{cases} \cos(\xi_0/2) &= \cos(\xi_2/2)\cos(\xi_1/2) - \sin(\xi_2/2)\sin(\xi_1/2)(\vec{n}_2 \cdot \vec{n}_1), \\ \sin(\xi_0/2)\vec{n}_0 &= \cos(\xi_1/2)\sin(\xi_2/2)\vec{n}_2 + \cos(\xi_2/2)\sin(\xi_1/2)\vec{n}_1 + \sin(\xi_2/2)\sin(\xi_1/2)(\vec{n}_2 \times \vec{n}_1), \end{cases}$$

and

$$\begin{aligned} \cos(\xi_0'/2) &= \cos(\xi_1/2)\cos(\xi_2/2) - \sin(\xi_1/2)\sin(\xi_2/2)\left(R_{(\vec{n}_2,\xi_2)}(\vec{n}_1)\cdot\vec{n}_2\right) \\ \sin(\xi_0'/2)\vec{n}_0' &= \cos(\xi_2/2)\sin(\xi_1/2)R_{(\vec{n}_2,\xi_2)}(\vec{n}_1) + \cos(\xi_1/2)\sin(\xi_2/2)\vec{n}_2 \\ &+ \sin(\xi_1/2)\sin(\xi_2/2)\left(R_{(\vec{n}_2,\xi_2)}(\vec{n}_1)\times\vec{n}_2\right). \end{aligned}$$

Since

$$\mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1) = \cos(\xi_2)\vec{n}_1 + \sin(\xi_2)(\vec{n}_2 \times \vec{n}_1) + (1 - \cos(\xi_2))(\vec{n}_2 \cdot \vec{n}_1)\vec{n}_2,$$

we have

$$\begin{cases} R_{(\vec{n}_2,\xi_2)}(\vec{n}_1) \cdot \vec{n}_2 = \vec{n}_1 \cdot \vec{n}_2, \\ R_{(\vec{n}_2,\xi_2)}(\vec{n}_1) \times \vec{n}_2 = \sin(\xi_2) \left( \vec{n}_1 - \left( \vec{n}_2 \cdot \vec{n}_1 \right) \vec{n}_2 \right) - \cos(\xi_2) \left( \vec{n}_2 \times \vec{n}_1 \right). \end{cases}$$

So we get the first result by direct substitution and simplification. The second result follows directly from the first one.  $\Box$ 

As a consequence, we have the following results.

**Corollary 4.** For rotations in  $\mathbb{R}^3$  we have

$$\mathbf{R}_{(\vec{n}_2,\xi_2)}\mathbf{R}_{(\vec{n}_1,\xi_1)} = \mathbf{R}_{\mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1)}(\xi_1)\mathbf{R}_{(\vec{n}_2,\xi_2)}$$

or

$$\mathbf{R}_{(\vec{n}_2,\xi_2)}\mathbf{R}_{(\vec{n}_1,\xi_1)}\mathbf{R}_{(\vec{n}_2,-\xi_2)} = \mathbf{R}_{\mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1)}(\xi_1)$$

**Corollary 5.** Let  $\xi \in (-\pi, \pi]$  and  $\vec{n} \in \mathbb{R}^3$  be given. Let  $\vec{n}'_0 = \vec{n}_0 = \vec{n}$ , and

$$(\vec{n}_l,\xi_l)$$
 is given, and  $\vec{n}'_l = R_{(\vec{n}_l,\xi_l)}(\vec{n}'_{l-1})$  for  $l = 1, 2, 3, \dots$ 

Then

$$\widehat{\operatorname{rot}}_{\vec{n}_l'}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_l}(\xi_l) \cdots \widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1) \widehat{\operatorname{rot}}_{\vec{n}_0}(\xi) \widehat{\operatorname{rot}}_{\vec{n}_1}(-\xi_1) \cdots \widehat{\operatorname{rot}}_{\vec{n}_l}(-\xi_l)$$

**Proof.** We have

$$\begin{aligned} \widehat{\operatorname{rot}}_{\vec{n}_{l}'}(\xi) &= \widehat{\operatorname{rot}}_{\mathbf{R}_{(\vec{n}_{l},\xi_{l})}(\vec{n}_{l-1}')}(\xi) \\ &= \widehat{\operatorname{rot}}_{\vec{n}_{l}}(\xi_{l})\widehat{\operatorname{rot}}_{\vec{n}_{l-1}'}(\xi)\widehat{\operatorname{rot}}_{\vec{n}_{l}}(-\xi_{l}) \\ &\vdots \\ &= \widehat{\operatorname{rot}}_{\vec{n}_{l}}(\xi_{l})\cdots\widehat{\operatorname{rot}}_{\vec{n}_{1}}(\xi_{1})\widehat{\operatorname{rot}}_{\vec{n}_{0}}(\xi)\widehat{\operatorname{rot}}_{\vec{n}_{1}}(-\xi_{1})\cdots\widehat{\operatorname{rot}}_{\vec{n}_{l}}(-\xi_{l}). \end{aligned}$$

**Example 1.** [1,2,6] Consider Euler angles, so  $\vec{n} = (\sin(\theta)\cos(\varphi), \sin(\theta)\sin(\varphi), \cos(\theta))$ . We verify directly that

$$\vec{n} = \mathbf{R}_{(\vec{e}_z, \varphi)} \left( \mathbf{R}_{(\vec{e}_y, \theta)}(\vec{e}_z) \right)$$

、

So, it follows that

$$\begin{split} \widehat{\operatorname{rot}}_{\vec{n}}(\xi) &= \widehat{\operatorname{rot}}_{\mathsf{R}_{(\vec{e}_{z},\varphi)}\left(\mathsf{R}_{(\vec{e}_{y},\theta)}(\vec{e}_{z})\right)}(\xi) \\ &= \widehat{\operatorname{rot}}_{\vec{e}_{z}}(\varphi)\widehat{\operatorname{rot}}_{\mathsf{R}_{(\vec{e}_{y},\theta)}(\vec{e}_{z})}(\xi)\widehat{\operatorname{rot}}_{\vec{e}_{z}}(-\varphi) \\ &= \widehat{\operatorname{rot}}_{\vec{e}_{z}}(\varphi)\widehat{\operatorname{rot}}_{\vec{e}_{u}}(\theta)\widehat{\operatorname{rot}}_{\vec{e}_{z}}(\xi)\widehat{\operatorname{rot}}_{\vec{e}_{u}}(-\theta)\widehat{\operatorname{rot}}_{\vec{e}_{z}}(-\varphi). \end{split}$$

# 11. Decomposition of rotation

The decomposition, or splitting, of a rotation into a sequence of two or more rotations is an important problem in mechanics, associated to robotics in  $\mathbb{R}^3$  [5,13], and in quantum computation, associated to gates in  $\mathbb{C}^2$  [6]. Thanks to the correspondence between rotations in  $\mathbb{R}^3$  and rotations in  $\mathbb{C}^2$ , we present in this section some recent decompositions of rotations in  $\mathbb{C}^2$  whose proof are based on decompositions results for rotations in  $\mathbb{R}^3$  presented in [4].

### 11.1. Computation of angle

We will use the trigonometric function  $atan2(\cdot, \cdot)$  to evaluate angles of rotation. This function is defined at every point  $(x, y) \neq (0, 0)$  in the plane  $\mathbb{R}^2$  [14]. Its value is uniquely determined by

$$\xi = \operatorname{atan2}(y, x) \text{ iff } \begin{cases} \xi \in (-\pi, \pi], \\ \cos(\xi) = x/\sqrt{x^2 + y^2}, \\ \sin(\xi) = y/\sqrt{x^2 + y^2}. \end{cases}$$

#### 11.2. Decomposition into two rotations

The first decomposition problem that we consider is the decomposition of a rotation as the product of two rotations. Two cases are considered. In the first case, we look for two axes for half-turn rotation. In the second case, the axes of rotation are given and we look for the angles of rotation.

11.2.1. Given angle : half-turn

For a unit vector  $\vec{n}$  and an angle  $\xi \in (-\pi, \pi]$  given, we would like to find two unit vectors  $\vec{n}_l$  and  $\vec{n}_r$  such that

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_l}(\pi) \widehat{\operatorname{rot}}_{\vec{n}_r}(\pi)$$

This result is used in [13].

For arbitrary  $\vec{n}_l$  and  $\vec{n}_r$ , and for  $\xi_l = \pi = \xi_r$ , the product of the two rotations leads to

$$\begin{cases} \cos(\xi/2) &= \cos^2(\pi/2) - \sin^2(\pi/2) \left( \vec{n}_l \cdot \vec{n}_r \right) = -\vec{n}_l \cdot \vec{n}_r, \\ \sin(\xi/2)\vec{n} &= \cos(\pi/2) \sin(\pi/2)\vec{n}_l + \cos(\pi/2) \sin(\pi/2)\vec{n}_r + \sin^2(\pi/2) \left( \vec{n}_l \times \vec{n}_r \right) = \vec{n}_l \times \vec{n}_r \end{cases}$$

so  $\vec{n}_l$  and  $\vec{n}_r$  are orthogonal to  $\vec{n}$ . We can select any unit vector  $\vec{n}_r$  orthogonal to  $\vec{n}$  and set  $\vec{n}_l = -R_{(\vec{n},\xi/2)}(\vec{n}_r) = R_{(\vec{n},\xi/2)}(-\vec{n}_r)$ , which is the same thing as selecting any unit vector  $\vec{n}_l$  orthogonal to  $\vec{n}$  and set  $\vec{n}_r = -R_{(\vec{n},-\xi/2)}(\vec{n}_l) = R_{(\vec{n},-\xi/2)}(-\vec{n}_l)$ .

11.2.2. Given axes

This problem is addressed in [4,5].

**Theorem 5.** Let a unit vector  $\vec{n}$  and an angle  $\xi \in (-\pi, \pi]$  be given. For two given unit vectors  $\vec{n}_l$ , and  $\vec{n}_r$  such that  $\vec{n}_l \times \vec{n}_r \neq \vec{0}$ , the following two statements are equivalent:

(i) there exist two angles  $\xi_l$  and  $\xi_r \in (-\pi, \pi]$  such that

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_l}(\xi_l) \widehat{\operatorname{rot}}_{\vec{n}_r}(\xi_r),$$

or

$$\mathbf{R}_{(\vec{n},\xi)} = \mathbf{R}_{(\vec{n}_l,\xi_l)} \mathbf{R}_{(\vec{n}_r,\xi_r)},$$

(ii)  $\vec{n}_l \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_r) = \vec{n}_l \cdot \vec{n}_r$ . Moreover •  $\xi_l = \operatorname{atan2}(b_l, a_l)$ , where

$$\begin{cases} a_l = (\vec{n}_l \times \vec{n}_r) \cdot (\vec{n}_l \times \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_r)), \\ b_l = (\vec{n}_l \times \vec{n}_r) \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_r). \end{cases}$$

•  $\xi_r = \operatorname{atan2}(b_r, a_r)$  where

$$\begin{cases} a_r = (\vec{n}_l \times \mathbf{R}_{(\vec{n},\theta)}(\vec{n}_r)) \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_l \times \vec{n}_r), \\ b_r = \vec{n}_l \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_l \times \vec{n}_r). \end{cases}$$

### 11.3. Decomposition into three rotations

Now we consider the following more complex problem studied in [4,5] for rotations in  $\mathbb{R}^3$ .

### 11.3.1. Statement of the general problem

**Problem 6.** Let a unit vector  $\vec{n} \in \mathbb{R}^3$  and an angle  $\xi \in (-\pi, \pi]$  be given. Let us consider three arbitrary given unit vectors  $\vec{n}_1$ ,  $\vec{n}_2$ , and  $\vec{n}_3$  in  $\mathbb{R}^3$ . Does there exist three angles  $\xi_1$ ,  $\xi_2$ , and  $\xi_3$ , such that

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_3}(\xi_3)\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1),$$

or, equivalently,

$$\mathbf{R}_{(\vec{n},\xi)} = \mathbf{R}_{(\vec{n}_3,\xi_3)} \mathbf{R}_{(\vec{n}_2,\xi_2)} \mathbf{R}_{(\vec{n}_1,\xi_1)}$$

This problem is solved by first considering conditions for the existence of  $\xi_2$ , and then we determine  $\xi_1$  and  $\xi_3$ .

**Looking for**  $\xi_2$ **.** We first observe that

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_3,\xi_3)} \mathbf{R}_{(\vec{n}_2,\xi_2)} \mathbf{R}_{(\vec{n}_1,\xi_1)}(\vec{n}_1) = \vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1).$$

So if the problem is solvable, we will have

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1) = \vec{n}_3 \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1)$$

The right-hand side is a quantity fixed by the data of the problem, so let us set

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1) = V \in [-1,1].$$

**Theorem 7.** Let  $V \in [-1,1]$  be a given real number. There exits  $\xi_2 \in (-\pi,\pi]$  such that

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_2, \tilde{\zeta}_2)}(\vec{n}_1) = V,$$

if and only if

$$|V - (\vec{n}_2 \cdot \vec{n}_3)(\vec{n}_2 \cdot \vec{n}_1)| \le \sqrt{\left(1 - (\vec{n}_2 \cdot \vec{n}_3)^2\right)\left(1 - (\vec{n}_2 \cdot \vec{n}_1)^2\right)}.$$

Moreover, let

$$L = \sqrt{\left( (\vec{n}_2 \times \vec{n}_3) \cdot (\vec{n}_2 \times \vec{n}_1) \right)^2 + \left( \vec{n}_2 \cdot (\vec{n}_1 \times \vec{n}_3) \right)^2},$$

and set

$$\begin{cases} A = (\vec{n}_2 \times \vec{n}_3) \cdot (\vec{n}_2 \times \vec{n}_1), \\ B = \vec{n}_2 \cdot (\vec{n}_1 \times \vec{n}_3), \\ C = (V - (\vec{n}_2 \cdot \vec{n}_3)(\vec{n}_2 \cdot \vec{n}_1)), \end{cases} and \begin{cases} a = A/L, \\ b = B/L, \\ c = C/L. \end{cases}$$

Using

$$\alpha = \operatorname{atan2}(b, a) = \operatorname{atan2}(B, A) \in (-\pi, \pi],$$

and

$$\beta = \operatorname{atan2}(\sqrt{1-c^2}, c) = \operatorname{atan2}(\sqrt{L^2 - C^2}, C) \in [0, \pi]$$

we get

$$\xi_2 = \alpha \pm \beta.$$

**Looking for**  $\xi_1$  and  $\xi_3$ . To determine  $\xi_1$  and  $\xi_3$  we have to consider two situations:  $R_{(\vec{n},\xi)}(\vec{n}_1) \neq \pm \vec{n}_3$  or  $R_{(\vec{n},\xi)}(\vec{n}_1) = \pm \vec{n}_3$ .

**Theorem 8.** Let us assume that there exists  $\xi_2$  such that

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1) = \vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1).$$

If  $\mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1) \neq \pm \vec{n}_3$  then there exists unique  $\xi_1$  and  $\xi_3$  in  $(-\pi,\pi]$  such that

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_3}(\xi_3)\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1).$$

or, equivalently

$$\mathbf{R}_{(\vec{n},\xi)} = \mathbf{R}_{(\vec{n}_3,\xi_3)} \mathbf{R}_{(\vec{n}_2,\xi_2)} \mathbf{R}_{(\vec{n}_1,\xi_1)}$$

•  $\xi_1$  is given by  $\xi_1 = \operatorname{atan2}(b_1, a_1)$  where

$$\begin{cases} a_1 = (\mathbf{R}_{(\vec{n},-\xi)}(\vec{n}_3) \times \vec{n}_1) \cdot (\mathbf{R}_{(\vec{n}_2,-\xi_2)}(\vec{n}_3) \times \vec{n}_1), \\ b_1 = \vec{n}_1 \cdot (\mathbf{R}_{(\vec{n},-\xi)}(\vec{n}_3) \times \mathbf{R}_{(\vec{n}_2,-\xi_2)}(\vec{n}_3)). \end{cases}$$

•  $\xi_3$  is given  $\xi_3 = \operatorname{atan2}(b_3, a_3)$ , where

$$\begin{cases} a_3 = (\mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1) \times \vec{n}_3) \cdot (\mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1) \times \vec{n}_3), \\ b_3 = \vec{n}_3 \cdot (\mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1) \times \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1)). \end{cases}$$

The next result is the equivalent Gimbal lock problem in  $\mathbb{R}^3$  for rotations in  $\mathbb{C}^2$ .

**Theorem 9.** Let us assume that there exists  $\xi_2$  such that

$$\vec{n}_3 \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1) = \vec{n}_3 \cdot \mathbf{R}_{(\vec{n}_2,\xi_2)}(\vec{n}_1)$$

*If*  $R_{(\vec{n},\xi)}(\vec{n}_1) = \pm \vec{n}_3$  then there exists a unique  $\delta_{\xi}$  such that for any  $\xi_1$  and  $\xi_3$  for which

$$\widehat{\operatorname{rot}}_{\vec{n}}(\xi) = \widehat{\operatorname{rot}}_{\vec{n}_3}(\xi_3)\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1)$$

or, equivalently

$$\mathbf{R}_{(\vec{n},\xi)} = \mathbf{R}_{(\vec{n}_3,\xi_3)} \mathbf{R}_{(\vec{n}_2,\xi_2)} \mathbf{R}_{(\vec{n}_1,\xi_1)},$$

we have  $\xi_1 \pm \xi_3 = \delta_{\xi}$  where  $\delta_{\xi} = \operatorname{atan2}(b, a)$  and

$$\begin{cases} a = (\vec{n}_2 \times \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_1)) \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_2 \times \vec{n}_1), \\ b = \vec{n}_2 \cdot \mathbf{R}_{(\vec{n},\xi)}(\vec{n}_2 \times \vec{n}_1). \end{cases}$$

11.3.2. Orthogonal axes of rotation

Under an orthogonality condition the problem is always solvable.

**Theorem 10.** If  $\vec{n}_1$  and  $\vec{n}_3$  are orthogonal to  $\vec{n}_2$ , then the necessary and sufficient condition of Theorem 7 holds true.

## 11.4. A mysterious looking example

The next result, called "mysterious looking theorem" in [1] for the special case ( $\vec{n}_1 = \vec{n}_3 = \vec{e}_z$ ,  $\vec{n}_2 = \vec{e}_y$ , and  $\vec{w} = \vec{e}_x$ ), is a key result for the construction of controlled multi-qubit gates in quantum computation [1]. It

is related to closed rotation sequences [15].

**Theorem 11.** Let us suppose given four angles of rotation  $\theta_l \in (-\pi, \pi]$  (l = 0, 1, 2, 3), and their four corresponding axes of rotation  $\vec{n}_l$  (l = 0, 1, 2, 3) such that

$$\widehat{\operatorname{rot}}_{\vec{n}_0}(\xi_0) = \widehat{\operatorname{rot}}_{\vec{n}_3}(\xi_3)\widehat{\operatorname{rot}}_{\vec{n}_2}(\xi_2)\widehat{\operatorname{rot}}_{\vec{n}_1}(\xi_1).$$

If  $\vec{n}_1 = \vec{n}_3$ , then there exist  $\hat{A}$ ,  $\hat{B}$ , and  $\hat{C}$ , such that

$$\hat{A}\hat{B}\hat{C}=\hat{\sigma}_{o},$$

and, for any unit vector  $\vec{w}$  orthogonal to  $\vec{n}_l$  (l = 1, 2, 3), we have

$$\widehat{\operatorname{rot}}_{\vec{n}_0}(\xi_0) = \widehat{\operatorname{Arot}}_{\vec{w}}(\pi)\widehat{\operatorname{Brot}}_{\vec{w}}(\pi)\widehat{\operatorname{C}}$$

Proof. Let us set

$$\begin{cases} \hat{A} = \widehat{\text{rot}}_{\vec{n}_{3}}(\xi_{3})\widehat{\text{rot}}_{\vec{n}_{2}}(\xi_{2}/2), \\ \hat{B} = \widehat{\text{rot}}_{\vec{n}_{2}}(-\xi_{2}/2)\widehat{\text{rot}}_{\vec{n}_{1}}(-\frac{\xi_{1}+\xi_{3}}{2}), \\ \hat{C} = \widehat{\text{rot}}_{\vec{n}_{1}}(\frac{\xi_{1}-\xi_{3}}{2}), \end{cases}$$

then

$$\hat{A}\hat{B}\hat{C} = \widehat{\text{rot}}_{\vec{n}_3}(\xi_3)\widehat{\text{rot}}_{\vec{n}_1}(-\xi_3)$$

So

$$\hat{A}\hat{B}\hat{C}=\hat{\sigma}_{o}$$

if and only if  $\vec{n}_1 = \vec{n}_3$  for arbitrary angle  $\xi_3$ . Moreover, for any unit vector  $\vec{w}$  orthogonal to  $\vec{n}_l$  (l = 1, 2, 3), we have

$$\widehat{\operatorname{rot}}_{\vec{w}}(\pi)(\vec{n}_i) = -\vec{n}_i \quad \text{for} \quad l = 1, 2, 3$$

So

$$\begin{split} \hat{A}\hat{\sigma}_{x}\hat{B}\hat{\sigma}_{x}\hat{C} =&\widehat{\mathrm{rot}}_{\vec{n}_{3}}(\xi_{3})\widehat{\mathrm{rot}}_{\vec{n}_{2}}(\xi_{2}/2)\widehat{\mathrm{rot}}_{\vec{w}}(\pi)\widehat{\mathrm{rot}}_{\vec{n}_{2}}(-\xi_{2}/2)\widehat{\mathrm{rot}}_{\vec{n}_{1}}(-\frac{\xi_{1}+\xi_{3}}{2})\widehat{\mathrm{rot}}_{\vec{w}}(\pi)\widehat{\mathrm{rot}}_{\vec{n}_{1}}(\frac{\xi_{1}-\xi_{3}}{2}) \\ =&\widehat{\mathrm{rot}}_{\vec{n}_{3}}(\xi_{3})\widehat{\mathrm{rot}}_{\vec{n}_{2}}(\xi_{2}/2)\widehat{\mathrm{rot}}_{\widehat{\mathrm{rot}}_{\vec{w}}(\pi)(\vec{n}_{2})}(-\xi_{2}/2)\widehat{\mathrm{rot}}_{\widehat{\mathrm{rot}}_{\vec{w}}(\pi)(\vec{n}_{1})}(-\frac{\xi_{1}+\xi_{3}}{2})\widehat{\mathrm{rot}}_{\vec{w}}(2\pi)\widehat{\mathrm{rot}}_{\vec{n}_{1}}(\frac{\xi_{1}-\xi_{3}}{2}) \\ =&\widehat{\mathrm{rot}}_{\vec{n}_{3}}(\xi_{3})\widehat{\mathrm{rot}}_{\vec{n}_{2}}(\xi_{2}/2)\widehat{\mathrm{rot}}_{-\vec{n}_{2}}(-\xi_{2}/2)\widehat{\mathrm{rot}}_{-\vec{n}_{1}}(-\frac{\xi_{1}+\xi_{3}}{2})\widehat{\mathrm{rot}}_{\vec{n}_{1}}(\frac{\xi_{1}-\xi_{3}}{2}) \\ =&\widehat{\mathrm{rot}}_{\vec{n}_{3}}(\xi_{3})\widehat{\mathrm{rot}}_{\vec{n}_{2}}(\xi_{2})\widehat{\mathrm{rot}}_{\vec{n}_{1}}(\xi_{1}) \\ =&\widehat{\mathrm{rot}}_{\vec{n}_{0}}(\xi_{0}). \end{split}$$

## 12. Conclusion

In this introduction to the mathematics of quantum physics and quantum computation, we put emphasis on the basic mathematical aspects of definition and operations on qubits. We started by a comprehensive introduction of a qubit as a unit element of  $\mathbb{C}^2$ , and its representations on spheres in  $\mathbb{R}^3$ . This introduction lead to the interpretation of Pauli operators as basic rotations in  $\mathbb{R}^3$ . Then we studied unitary operators, and their link to rotations in  $\mathbb{R}^3$  has been established with the density operator associated to a qubit. Formulas for the composition and non-commutativity of two rotations have been established. We completed this paper by some problems of decomposition, or splitting, of unitary operators on  $\mathbb{C}^2$ , and rotations in  $\mathbb{R}^3$ , which are related to gates for quantum computation.

Conflicts of Interest: The author declares no conflict of interests.

**Funding Information:** This work has been financially supported by an individual discovery grant from the Natural Sciences and Engineering Research Council of Canada.

#### References

- Nielsen, M.A., & and Chuang, I.L. (2010). *Quantum Computation and Quantum Information*. 10th Anniversary Edition, Cambridge University Press.
- [2] Olivares, O. (2021), Lecture Notes on Quantum Computing. University of Milan, Italy.
- [3] Yepez, J. (2013). *Lecture Notes: Qubit Representations and Rotations*. Physics 711, Topics in particles & fields, University of Hawaï at Manoa, USA.
- [4] Dubeau, F. (2025). Composition, non-commutativity, and decompositions of finite rotations, *Engineering Reports*, 7, e13107, 1-14.
- [5] Piovan, P., & Bullo, F. (2012). On Coordinate-Free Rotation Decomposition: Euler Angles about Arbitrary Axes. IEEE Transactions on Robotics, 28, 728-733.
- [6] McMahon, D. (2007). Quantum Computing Explained. Wiley-Interscience, John Wiley & Sons Inc.
- [7] Halmos, P.R. (1958, 1974). *Finite-Dimensional Vector Spaces*. Litton Educational Publishing Inc, and Springer-Verlag New York Inc.
- [8] Hsu, H.P. (1969). Vector Analysis. Simon & Schuster, New York, USA.
- [9] Jones, F. (2004). Text for Honors Calculus III/IV, Rice University, USA.
- [10] Cheng, H., & Gupta, K.C. (1989). An historical note on finite rotations. Journal of Applied Mechanics, 56, 139-145.
- [11] Paul, B. (1963). On the composition of finite rotations. The American Mathematical Monthly, 70(8), 859-862.
- [12] Choudhury, A.A., Keil, M.J., & Rodriguez, J. (2008). On the commutativity of sequential finite rotations. *International Journal of Mechanical Engineering Education*, 36(2), 113-119.
- [13] Wohlhart, K. (1992). Decomposition of a finite rotation into three consecutive rotations about given axes. *Proceedings* of VI-th International conference on Theory of Machines and Mechanics, Czechoslovakia, Liberec, 325-332.
- [14] Wikipedia. atan2(y,x) formula. https://en.wikipedia.org/wiki/Atan2.
- [15] Bhat, S.P., & Crasta, N. (2015). Closed Rotation Sequences. Discrete & Computational Geometry, 53, 366–396.



© 2025 by the authors; licensee PSRP, Lahore, Pakistan. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (http://creativecommons.org/licenses/by/4.0/).